

KPL DAB Reagent Set

Catalog No.
5510-0031 (54-10-00)

Size
500 Slides

DESCRIPTION

DAB (3,3'-diaminobenzidine) deposits a brown specific stain in the presence of horseradish peroxidase (HRP). The substrate is useful for immunohistochemical and immunoblotting applications.

CONTENTS

The KPL DAB Reagent Set provides a three component liquid substrate system in convenient dropper bottles:

15 mL KPL Tris Buffer (0.1 M)
10 mL KPL DAB Solution (25 mg/ml)
10 mL KPL Peroxide Solution (0.5%)

Sufficient reagents are supplied to stain approximately 500 slides.

STORAGE/STABILITY

Store reagents at 2-8°C. Stable for a minimum of one year from date of receipt when stored at 2-8°C.

SUGGESTED REAGENTS NOT INCLUDED

1. Primary antibody.
2. KPL Peroxidase blocking solution or H₂O₂.
3. HRP-labeled secondary antibody or streptavidin.
4. KPL Contrast BLUE or hematoxylin.
5. Isopropyl alcohol.
6. Mounting media.
7. 0.1 M Tris-HCl or PBS
(See SOLUTION PREPARATION).

PREPARATION

Note: Warm reagents to room temperature before use.

1. Add 3 drops (150 µL) KPL Tris Buffer Concentrate to 5 mL reagent quality water.
2. Add 2 drops (100 µL) KPL DAB Solution.
3. Add 2 drops (100 µL) KPL Peroxide Solution.

Mix solution thoroughly and use immediately. The working substrate solution contains 0.5 mg/ml DAB and 0.1% H₂O₂.

STAINING PROCEDURE

1. Rehydrate paraffin embedded sections through graded alcohol (3 minutes each in 100%, 80%, 40%, and 20% EtOH) to water. Other samples listed below do not require rehydration. Frozen sections must be thoroughly dried before use.
2. Block endogenous peroxidase activity by immersing samples in KPL Peroxidase Blocking Solution as follows (If using H₂O₂ see TROUBLESHOOTING):
 - a. Frozen sections - 45 seconds
 - b. Paraffin sections - 4 minutes
 - c. Cytospin preparations - 45 seconds
 - d. Blood films - 45 seconds
 - e. Touch or squash preparations - 1 minute
 - f. Floating or whole sections - 5 minutes
3. Rinse five minutes in reagent quality water.
4. Soak in 0.1 M Tris-HCl or PBS 10 minutes.
5. Treat sample with primary antibody diluted in Tris-HCl or PBS 15-20 minutes.

NOTE: Extended incubation may improve sensitivity.

6. Wash sample with Tris-HCl or PBS 10 minutes.
7. Incubate sample with biotinylated antibody, directed against the primary antibody host species, 15-20 minutes. If using HRP-labeled secondary antibody, go to step 9.
8. Wash as in Step 6.
9. Shake off excess buffer and incubate sample with HRP Streptavidin or HRP-labeled secondary antibody diluted in Tris-HCl or PBS, 15-20 minutes.
10. Wash as in Step 6. (Prepare KPL DAB substrate during this step).
11. Shake off excess buffer and cover section with KPL DAB substrate.
12. Incubate 10 minutes at room temperature out of direct light.
13. Rinse slide 2-3 minutes in reagent quality water.
14. Counterstain with KPL Contrast BLUE, or hematoxylin, if desired:
 - a. Paraffin embedded and frozen sections for 3 minutes.
 - b. Touch preparations, cytospin preparations and blood films for 30-45 seconds.

KPL DAB Reagent Set

Catalog No.

5510-0031 (54-10-00)

Size

500 Slides

15. Rinse thoroughly in 2-3 changes of isopropyl alcohol or until excess stain is removed from slide. DO NOT USE WATER OR ETHANOL SOLUTIONS.
16. Air dry and mount with aqueous or xylene-based mounting medium.

Note: KPL TrueBlue™ rPeroxidase Substrate is recommended for double labeling procedures.

RESULTS

1. Sites of enzyme activity range from light to dark brown.
2. If counterstained, nuclei appear a contrasting blue.
3. Sections not reacted with primary antibody as a negative control should not develop a brown tint.
4. To prevent background, further dilution of primary antibody or HRP-labeled reagent may be required.

TROUBLESHOOTING

1. Always incorporate appropriate positive and negative controls.
2. Instant development of brown color indicates that the primary antibody or peroxidase-labeled reagent must be further diluted.
3. Prolonged incubation in substrate may increase background and inhibit nuclear counterstaining.
4. As an alternative method to block endogenous peroxidase, incubate slides for 30 minutes in 0.3% (w/v) H₂O₂ in absolute methanol followed by a 10-15 minute rinse in 0.1 M Tris-HCl, pH 7.6 or PBS.

DISPOSAL

The following method of disposal is recommended for solutions containing DAB:

1. Add 100 mL of household bleach to 2 liters of water. Pour solution into a 1 gallon plastic bottle.
2. Pour waste DAB solution into the bleach solution and mix by shaking. No more than 500 mL of DAB solution should be added.
3. After last addition, allow container to stand at least 24 hours before discarding.

BUFFER PREPARATION

0.1 M TRIS-HCl

1. Dissolve 121 g Tris in 500 mL reagent quality water.
2. Adjust pH to 7.6 with 2 M HCl (approximately 300 mL).
3. QS to 1 L with reagent quality water to obtain a 1 M stock.
4. Dilute 1 part stock from step 3 with 9 parts reagent quality water and mix well.

Phosphate Buffered Saline (PBS)

1. Add PBS (0.01 M), 8.0 g NaCl, 0.2 g KCl, 1.44 g Na₂HPO₄, 0.24 g KH₂PO₄.
2. Adjust pH to 7.4 with 2 M HCl.
3. QS to 1 L with reagent quality water and mix well.

REFERENCES

1. Nakane, P.K. and Pierce, G.B. Jr. (1966). J. Histochem. Cytochem. 14: 929.
2. Sternberger, L.A. et. al. (1970). J. Histochem. Cytochem. 18: 315.
3. Hsu, S.M. and Ree, H.J. (1980). Am. J. Clin. Pathol. 74:32.
4. DeJong, A. et. al. (1985). Histochem. 17: 1119.
5. Graham, R. et. al. (1965). J. Histochem. Cytochem. 13:150.
6. Hanker, J.S. et. al. (1977). Histochem. 9: 789.
7. Nakane, P.K. (1968). J. Histochem. Cytochem. 16: 557.
8. Graham, R.C. Jr. and Karnovsky, M.J. (1966). J. Histochem. Cytochem. 14: 291.
9. Hanker, J.S. (1977). Osmophilic Reagents in Electronmicroscopic Histochemistry in Progress in Histochemistry and Cytochemistry, VCH Publishers, Deerfield Beach, FL.
11. Kelly, J. et. al. (1987). J. Immunol. Meth. 96: 127.
12. Segasothy, M. et. al. (1988). Am. J. Clin. Pathol. 90: 691.
13. Molin, S.O. et. al. (1978). J. Histochem. Cytochem. 26:412.
14. Pickel, V.T. et. al. (1976). J. Histochem. Cytochem. 24:792.

KPL DAB Reagent Set

Catalog No.

5510-0031 (54-10-00)

Size

500 Slides

PRODUCT SAFETY AND HANDLING

This product is considered hazardous as defined by the Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, consult MSDS. Dispose of this product as instructed above.

RELATED PRODUCTS

CAT. NO.

KPL Blocking Solution Concentrate	5560-0006 (71-00-10)
KPL Contrast BLUE	5540-0002 (71-00-06)
KPL TrueBlue® HRP Substrate	5510-0049 (71-00-64)

PRINCIPLE

The application of antibodies and other proteins covalently coupled to horseradish peroxidase (HRP) in immunohistology is well documented ⁽¹⁻⁴⁾. It is the most frequently used label for immunohistologic techniques. In the presence of peroxide, HRP catalyzes the oxidation of phenols, naphthols, diamines, aminophenols, indophenols, etc., forming chromogenic products visible by light microscopy. Most commonly employed are 3-amino-9-ethylcarbazole ⁽⁵⁾, p-phenylenediamine/catechol ⁽⁶⁾, 4-chlorol-naphthol ⁽⁷⁾ and diaminobenzidine (DAB) ⁽⁸⁾. Although a suspected carcinogen, DAB is the most widely accepted donor substrate for peroxidase immunohistochemistry, since it provides a reaction product insoluble in alcohols and xylene.

The oxidation of DAB results in formation of a free radical intermediate which polymerizes to form a brown product. DAB may be employed for demonstration of endogenous peroxidase and catalase activity; cytochrome oxidase; cupric ferrocyanide; and hemoproteins such as hemoglobin, myoglobin, and cytochrome c. Treatment of the DAB product with osmium, silver, cobalt or nickel will intensify final color. Reaction with osmium tetroxide results in an electron opaque osmium black useful for ultrastructure research.

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis.